



PLATTE RIVER RECOVERY IMPLEMENTATION PROGRAM

Whooping Crane Monitoring Protocol – Migrational Habitat Use in the Central Platte River Valley

I. INTRODUCTION

In 2007, the Platte River Recovery Implementation Program (“Program” or “PRRIP”) began its 13-year First Increment and implementation of an Adaptive Management Plan (“AMP”) to learn more about the physical processes of the central Platte River and the response of four target species to management actions: interior least tern (*Sterna antillarum*), piping plover (*Charadrius melanotos*), whooping crane (*Grus americana*), and pallid sturgeon (*Scaphirhynchus albus*). The AMP includes several protocols for monitoring target species, habitat, and physical processes to better understand interrelationships and provide data for evaluating species response to management actions. This document serves as the whooping crane monitoring protocol for the Program.

Information from this protocol will be used to help evaluate the biological response of whooping cranes and habitat to the land and water management activities of the Program. Several critical scientific and technical uncertainties about Program target species, physical processes, and the response of the target species to management actions will be the focus of the application of rigorous adaptive management in the First Increment through implementation of the Program’s AMP. These uncertainties are captured in statements of broad hypotheses on pages 14-17 of the AMP and, as a means of better linking science learning to Program decision-making, those uncertainties comprise a set of “Big Questions” that provide a template for linking specific hypotheses and performance measures to management objectives and overall Program goals (see PRRIP 2007-2010 Synthesis Report, 2011) .

Two “Big Questions” relate directly to whooping cranes:

- **Big Question #5** – Do whooping cranes select suitable riverine roosting habitat in proportions equal to its availability?
- **Big Question #10** – How do Program management actions in the central Platte River contribute to least tern, piping plover, and whooping crane recovery?

These uncertainties led to the development of a specific management objective for the whooping crane and indicators related to that objective, as noted in the AMP:

| | |
|-----------------------------|--|
| Management Objective | <i>Contribute to the survival of whooping cranes during migration</i> |
| Indicators | <ul style="list-style-type: none">* Increase area of suitable roosting and foraging habitat* Increase crane use days* Increase proportion of whooping crane population use |

To assess progress toward this objective and learn about the major whooping crane uncertainties, several finer-scale priority hypotheses were developed by Program participants. In 2010, those hypotheses were sequenced to develop a smaller set of Tier 1 hypotheses to receive focused attention in the First Increment. For whooping cranes, those Tier 1 hypotheses are:

- **WC-1:** Whooping crane use will increase as a function of Program land and water management activities.
- **WC-3:** Whooping crane use is related to habitat suitability. Riverine habitat suitability for whooping cranes is a function of channel characteristics such as water depth, channel width, and unobstructed-view widths.

This monitoring protocol is intended to provide standard implementation guidance for collecting whooping crane use and habitat (i.e., landscape level attributes at roost sites and diurnal use sites) data necessary to test the Tier 1 whooping crane hypotheses, assess progress toward meeting the whooping crane management objective, and evaluate learning related to the whooping crane Big Questions. Use of the phrase “suitable channel habitat” relates to the Program’s established minimum habitat criteria (see



PRRIP Draft Whooping Crane Minimum Habitat Criteria (Updated 8-1-12)). This protocol will be used by the Program to gather information on whooping crane habitat use and to provide an index of whooping crane abundance in the study area. It is understood that regardless of survey method not all whooping cranes are certain of being detected during migration and therefore full implementation of this or any other protocol will not represent complete whooping crane use of the central Platte River valley.

II. PURPOSE

The purpose of this monitoring protocol is to describe the conceptual design, study methods, and procedures that will be used annually to gather repeatable information on whooping crane stopovers in the central Platte River valley, Nebraska. The protocol outlines information the Program's Whooping Crane Monitoring Consultant (Consultant) will collect in the field, as well as from FWS and state agencies, and describes the procedures to be used for these specific objectives:

- 1) Detect and confirm whooping crane stopovers in the study area (Appendix A) – systematic, but targeted aerial surveys of the river channel and wetlands within the study area will be conducted and the data will be used to comparatively evaluate changes in the frequency and distribution of stopovers within the study area over time. The Whooping Crane Tracking Partnership's (WCTP) telemetry project locations and opportunistic locates will also be used to detect whooping crane stopovers in the study area; however, telemetry data will only be used by the Consultant to relocate whooping cranes using the study area that were already observed by the monitoring crew. Additional whooping crane stopover locations identified via telemetry will be reported to the Consultant after the whooping crane group leaves the study area.
- 2) Landscape Data Collection – Basic landscape source data of whooping crane use-sites in the study area will be collected through this protocol. Habitat metrics will be collected at confirmed systematic, opportunistic, and telemetry-marked whooping crane use sites. This information will be used in future use/availability analyses using aerial photography, Geographic Information System (GIS) information, LiDAR data, and appropriate landscape data collected from other protocols. Currently the Program has available a complete land use/land cover GIS analyses of 1998 (baseline) and 2005 color infrared photography. Continued regular collection of landscape data sources of the study area through other protocols such as aerial photographs, LiDAR, geomorphology monitoring, GIS data, and annual habitat availability assessments will enable future habitat use/availability analyses.

III. DESIGN CONSIDERATIONS AND SPECIFICATIONS

III.A. Area of Interest

The area of interest for monitoring whooping crane migrational habitat use consists of an area 3.5-miles either side of the Platte River beginning at the junction of U.S. Highway 283 and Interstate 80 near Lexington, Nebraska and extending eastward to Chapman, Nebraska. When side channels of the Platte River extend beyond the 3.5-mile area, a 2-mile area is included around these channels (see Appendix A)).

III.B. Project Design

This protocol collects information on whooping cranes using the central Platte River, not necessarily on the entire whooping crane population. This may bias the sample for making inference to the entire whooping crane population. In addition, the results from this protocol may not be representative of the population, or subgroup of the population using the central Platte River, because of the use of multiple observations per crane group and/or the lack of use by unique crane groups in the analysis (i.e., pseudo-replication). Options for addressing pseudo-replication are discussed in Section IV.D.

III.B.1. Detecting/Locating Whooping Crane Stopovers

Whooping crane stopovers will be documented using systematic aerial surveys, telemetry locations, and opportunistic sighting reports. Crane groups detected with systematic aerial surveys will have probabilities of inclusion estimated for the sample while crane groups detected opportunistically will compromise a non-



probability based sample. Systematic riverine surveys cover the study area from east to west with equal effort, however, biases associated surveying the ‘main channel’ will be accounted for through detection trials. Inferences from analyses of riverine survey data will be applicable to all Platte River channels. Systematic return surveys cover the off-channel portion of the study area most likely to be used by whooping crane groups at or near sunrise. Targeted wetlands and all landcover within a ½ mile area of the return transect will be surveyed under this protocol. Inferences from analyses of return survey data are applicable to a ½ mile buffer surrounding the return transect. Inferences from analyses of opportunistic surveys will contain biases associated with unequal sampling effort that cannot be accounted for and therefore may not represent actual crane use of the study area. Telemetry data, public reports, and reports from other monitoring efforts in the valley will also be used to document occurrences of whooping crane stopovers in the study area. These sighting reports increase the opportunity to gather habitat use information.

Relative probabilities of detection and efficiency of sighting whooping crane groups using aerial surveys will be estimated through protocol implementation over the years (e.g., use of decoys, known birds in the area, etc.). As detection may differ across various habitat conditions surveyed (e.g., wide versus narrow channels, corn versus grassland, etc.), detection probabilities will be estimated via detection trials (i.e., decoys and/or telemetry-marked birds) and results will be used in analyses to account for potential biases.

Aerial Survey

Aerial surveys will be used to detect whooping crane use-site locations (i.e., roost locations) in the study area. Systematic, but targeted aerial surveys are necessary to develop information on the spatial and temporal distribution of whooping crane use site locations in and along the Platte River for comparative evaluations. The design of the aerial surveys is intended to maximize opportunities to observe whooping crane use sites throughout the study area. Daily flights will be conducted in early morning during the period when whooping cranes are most likely to be in route between the wintering and breeding grounds and at or near their roost location. Daily flights will take place over the main river channel (river transects) and upland areas with high densities of wetlands within the study area (return transects; Appendix B). The “main river channel” is defined as the widest channel when all channels have flowing water. It is recognized that this protocol only samples river channels and a 1-mile area surrounding the targeted wetland return transects and does not sample other areas of the study area (see inferences defined in III.B.1). River transects will be flown east to west and return transects will be flown west to east.

Ground Confirmation

The consultant will deploy at least 1 person on the ground daily to confirm sightings that cannot be confirmed from the air via photographs or positive identification. In the event more than 1 unconfirmed crane group is observed on the same day, one or more of the air crew members will return to the area the additional unconfirmed crane groups are located to attempt to confirm them as well. A minimum of 2 hours will be spent observing for crane groups to leave the roost to forage or to migrate. Once the crane group(s) are positively confirmed or 2 hours has been devoted to confirming the birds from the ground, monitoring will be complete for the day. All ground monitoring will be conducted from public roads unless the consultant has permission to enter a property.

Telemetry Locations

Locations of whooping crane use of the study area may be documented by the Whooping Crane Tracking Partnership. Once telemetry-marked whooping cranes leave the study area, information regarding use of the study area will be provided to the consultant to document all known whooping crane stopover occurrences in the database. All telemetry-marked whooping crane groups that use the study area will be included in detection trials. Similar to unmarked whooping cranes, the probability telemetry-marked whooping cranes are included in the systematic sample will be estimated for groups observed by systematic aerial surveys.



Opportunistic Locates

Birdwatchers, outdoor enthusiasts, farmers, and other survey efforts might make initial observations of whooping crane groups in the study area. Sighting reports from these and other groups (labeled “opportunistic locates”) may provide additional information on crane stopover occurrences, but the conclusions are only applicable to the areas searched by the people that would report a sighting. An analysis of habitat use by cranes sighted opportunistically is outlined in this protocol, however, locations of whooping cranes obtained through this method are biased and quantifying the bias due to the location and amount of effort expended to obtain these observations is not planned.

Survey Detection Rates

Telemetry-marked whooping cranes and/or decoys can be used to estimate the probability whooping crane groups are detected during aerial surveys. Decoys will be randomly placed on active Platte River channels and within ½ mile of the upland return transect. Decoys may be placed on private, Program, governmental or non-governmental organizations’ lands in which Program personnel (EDO staff, partners, etc.) have permission to access. Program personnel will place decoys within the same landcover class and as close to the randomly generated locations as possible (i.e., nearest location that conforms to Program Minimum Habitat Criteria) and will record the UTM location of the decoy. Aerial survey crews will not know the location of decoys while conducting the survey and detection rates will be calculated as the percentage of decoys observed. Detection trials will also be used in detectability models to account for unequal probabilities of inclusion in the sample associated with surveying the main channel and potential differences in detection within the various off-channel landcover classes.

III.C. Timing

Aerial surveys of the study area will be conducted in the spring from March 6 to April 29 and in the fall from October 9 to November 15. These dates generally encompass the fifth and ninety-fifth percentile of initial observation dates of whooping cranes in Nebraska, 2007–2016, as well as the scheduled monitoring seasons, spring 2001–fall 2016 (Appendix C & D). Opportunistic and telemetry locations will be documented throughout the year; however, data collection efforts described in this protocol may or may not be implemented outside the scheduled monitoring seasons.

IV. METHODS

IV.A. Definitions

Crane Group – one or more cranes in a migrating unit. The group may consist of an individual crane, a family unit, or small flock. The social make-up of crane groups should be recorded whenever possible.

Stopover – Use of the study area during spring or fall migration.

Use Location – A location of a crane group that occurs in any landcover class within the study area. A single crane group may, and likely will have more than one use location per day.

Use Site – A location of a crane group within a wetland landcover class (wetted channel, open water pit/pond/lake, etc.) in the study area. Use sites are a special type of use location in that they only occur in wetland habitat. Use sites will be assigned a *Use Site ID* (e.g., 01) as well as a *Location ID* (e.g., A). A single crane group may have more than one use site per day.

Sighting – observation of a crane group in the study area.

Confirmed Sighting - Observation made by a State or Federal biologist or officer or by other known qualified observer (trained ornithologist or birder with experience identifying of whooping cranes).

A photograph may also be used to confirm sightings. Aerial survey crew with previous aerial whooping crane observations may confirm a crane group during the survey.

Probable Sighting - No confirmation made by State or Federal biologist or officer or by other known qualified observer, yet details of the sighting seem to identify the birds as whooping cranes. To be classified as a probable sighting each of the following factors must be met: (1) location of sighting is



within normal migration corridor and is an appropriate site for whooping cranes; (2) date of sighting is within period of migration; (3) accurate physical description; (4) number of birds is reasonable; (5) behavior of the birds does not eliminate whooping cranes; and (6) good probability that the observer would provide a reliable report.

Unconfirmed Sighting - Sighting details meet some, but not all six factors listed for a probable sighting.

Systematic Sighting – Aerial observation of whooping cranes made during a scheduled survey along specified transects as well as ground observations that are the result of such an aerial survey. Systematic observations do not include aerial observations made outside the scheduled monitoring season or off scheduled flight paths.

Opportunistic Sighting – Aerial observation of whooping cranes not made during a scheduled survey along specified transects as well as ground observations that are not the result of a scheduled aerial survey along transects. Opportunistic sightings include observations made outside the scheduled monitoring season, off scheduled flight paths, and observations made as the result of public, telemetry, etc. reports.

IV.B. Field Techniques

IV.B.1. Detecting/Locating Whooping Crane Stopovers

Three methods will be used to locate migrating whooping crane stopovers along the central Platte River during spring and fall migration: aerial surveys, telemetry data, and opportunistic locates. The Program's Technical Advisory Committee may choose to implement each protocol component as necessary to obtain needed information, for example changing the survey effort based on results of past surveys.

Aerial Survey

Daily aerial surveys, weather permitting, will be conducted along the central Platte River valley between Lexington and Chapman, Nebraska to locate spring and fall migrating whooping crane groups. Aerial surveys will take place from March 6 to April 29 in the spring and October 9 to November 15 in the fall. These dates encompass the fifth and ninety-fifth percentile of initial sighting dates for all recorded sightings of whooping crane groups in Nebraska, 2001 – 2016 (timeframe the monitoring protocol has been implemented) as well as 1975 – 1999 (Jane Austin, USGS Northern Prairie Wildlife Research Center, pers. comm.). This protocol intends to collect a sample during possible migration time and does not intend to survey the entire time-period it would be possible for a crane group to migrate through the study area. Therefore, the survey dates will not be extended during times of delayed migration or initiated earlier in times of early migration. However, if the survey period extends past the migration time in a given season, the surveys will be stopped using the following rules. For the spring survey, flights will be discontinued 5 days after the last normally migrating whooping cranes have departed Aransas, if no whooping cranes have been sighted in the central Platte valley for 5 days, and there are no recent (5 days) reports of whooping cranes in the Central Flyway south of the Platte River. For the fall survey, flights will be discontinued if no whooping cranes have been sighted in the central Platte valley for 5 days, and there are no recent (5 days) reports of whooping cranes in the Central Flyway north of the Platte River. The Program Manager or Biologist responsible for managing these surveys will be in contact with Wade Harrell (or other Aransas official) at (361) 286-3559 to obtain information related to bird departure/arrival from Aransas and will determine whether to continue aerial surveys or not.

A Cessna 172 or similar aircraft will fly at a speed of 100 mph, as safety allows. One plane will fly the area between Chapman and the Nebraska Highway 10 (Minden) Bridge (the east leg). The second plane will fly the area between the Minden Bridge and the Lexington Bridge (the west leg). Two observers in addition to the pilot will be in each plane. Surveys will begin between a half-hour before sunrise and sunrise, unless weather during this time period precludes beginning the survey. All attempts should be made to begin the survey a half-hour before sunrise. If the survey cannot begin during this time period due to weather/visibility requirements, the survey start time will be extended up to two hours after sunrise. Surveys may be canceled two hours after sunrise due to unsafe weather conditions (e.g., rain, snow, fog, high winds) or if there is significant snow cover on the ground that greatly impedes the surveyors chances



of locating a whooping crane group. Cancelled flights will be entered into the database with an explanation of why the flight was cancelled (i.e., include weather conditions, etc.).

All aerial surveys will be flown such that the flight direction when flying the river transect will be away from the rising sun. To help address the concern that one end of the river transect will always be flown early and the other late, there will be two start locations for each leg (east side and west side) of the study area. For the eastern section, on day one the flight will begin at Chapman, fly the river west to Minden, fly the primary wetland return transect back to Chapman, and fly the secondary wetland return transect back to the Burlington Northern railroad near Grand Island. On day two the flight will begin at the Wood River Bridge, fly the river transect west to Minden, fly the primary wetland return transect back to Chapman, fly the rest of the river transect from Chapman to Wood River, and finish up on the secondary wetland return transect between Wood River and HWY 281 near Grand Island. For the western section, on day one the flight will begin at Minden, fly the river west to Lexington, and then fly the primary wetland return transect back to Minden. On day two the flight will begin at the Odessa Bridge, fly the river transect west to Lexington, fly the primary wetland return transect back to HWY 10 south of the Minden Bridge, fly the rest of the river transect from Minden to Odessa, and then fly the secondary wetland return transect from Elm Creek back to HWY 10 north of the Minden Bridge. These patterns will continue through the survey period.

During the river transect, observers will be situated such that the main channel(s) can be clearly viewed by both observers looking out the passenger side of the plane. This will necessitate that the plane fly just south of the main channel. An exception to this rule occurs west of the Overton Bridge when the plane will fly over Jeffery Island and observers position themselves to observe both the north and south channel of the Platte River. On the return transect, observers will look out different sides of the plane so that they can survey a half-mile on each side of the targeted wetland return transect. This design provides a systematic aerial survey to locate whooping crane groups within the channel as well as outside the channel within the targeted wetland survey area. If additional wetlands are created or are identified, the Program's Technical Advisory Committee may choose to alter the return transects to sample these areas and indices of use (e.g., # whooping cranes/flight mile) may need to be adjusted to account for change in area surveyed. Again, it is recognized that this sampling scheme over-samples the river and targeted wetland areas compared to upland areas not covered by the return transects.

All transects will be flown at 750' altitude unless FAA regulation or snow goose migration dictate a higher altitude (e.g., a minimum of 1000' altitude when flying over towns and cities). The 750' altitude for transects is selected for safety reasons. Extremely large numbers of migratory waterfowl are present in the central Platte River valley each spring so flights may be flown at 2,000' altitude when the safety of the flight crew is in risk. The 2,000' altitude allows pilots to fly over airborne waterfowl to decrease the chance of collision. If a suspected whooping crane is seen, the plane is encouraged to circle to an altitude of 500' (when safety allows) to provide a better viewing opportunity of the suspected whooping crane.

Each plane will have a global position system (GPS) unit to aid in flight-path orientation and documentation of crane group locations. UTM coordinates taken either from the plane's GPS system or hand held unit will be recorded on the data sheet. The aerial survey crew will photograph the whooping crane group and the general location using a digital camera with an 18 × 105 mm Vibration Reduction (VR) zoom lens or similar setup approved by the Program. All observations will be recorded on the aerial observation datasheet. Deviations from return transect flight paths may be implemented (and documented on the datasheet) to relocate or confirm crane groups in the area; however, river transects should be completed without interruption. If the crane group cannot be confirmed from the air, the plane will maintain visual contact with the crane group and direct a ground monitoring crew to the location to confirm the sighting.



Telemetry Location Data

The Program's Technical Point of Contact will not provide information about the use of the study area by radio-marked whooping cranes to the Consultant until they have first been observed and documented by the Consultant's crew.

Opportunistic Locates

The quality and timing of public sighting reports are highly variable. For example, several reports of a single group may be made by different individuals; sightings may be reported after the group has left the area; geese, white sandhill cranes, pelicans, or egrets may be reported as whooping cranes; etc. In an effort to document the validity of a sighting in a timely manner, a toll free number will be used to relay reports of possible whooping crane sightings to US Fish and Wildlife Service. This number should be publicized at local areas frequented by birders, FWS offices, NGPC offices, and possibly in newspapers, to mail carriers, bus drivers, etc. The monitoring crew will attempt to confirm all "probable" and "unconfirmed" sightings reported to be in the study area.

The monitoring crew will fill out ground monitoring observation forms for all effort expended to locate probable and unconfirmed sightings of crane groups in the study area. Information on all confirmed and probable sightings made by the monitoring crew will be forwarded to the FWS Nebraska field office. Incidental observations reported to the monitoring crew from inside or outside the study area will be forwarded to the FWS Nebraska field office, Whooping Crane Migration Information Coordinator.

Survey Detection Rates

Whooping crane decoys will be placed at randomly selected locations in the riverine and upland survey areas in which Program personnel has permission to access. Aerial crews will not be aware of the presence of the decoys during the flight. When the aerial crew observes a decoy, the location of the sighting should be relayed to the consultant's designated point of contact for confirmation of the decoy location. Decoy observations will be recorded on the aerial observation datasheet and information pertaining to detected and undetected decoys will be entered into the Program's electronic database. Detected and undetected telemetry-marked whooping cranes can also provide the Program information on aerial detection rates.

Aerial flight crews shall not be notified of the location of whooping cranes in the study area as this information clearly influences detection probabilities. In the event aerial flight crews are notified of the location of whooping cranes in the study area, they will include a note in the datasheet indicating they were aware of the presence of whooping cranes prior to the survey and will describe how they were made aware of this information.

IV.B.2 Crane Group Numbering

The *Crane Group ID* will consist of the following information: year; "SP" for spring monitoring period or "FA" for fall; sequential number (e.g. 2002FA01, 2002FA02, 2002FA03, etc.). Any time a crane group is observed in the study area by the survey crew, a new *Crane Group ID* will be assigned unless the surveyor notes on the data sheets the reasons why they believe this is a previously recorded group (using their professional judgment). In this case, the same *Crane Group ID* will be used. FWS crane group numbers for confirmed sightings will be included in the Program database and linked to the Program crane group numbers. This will assist in future cross-referencing between unique whooping crane groups in the FWS and Program databases.

Photographs taken of crane use-sites observed from the air will be used to locate use sites on Google Earth or other aerial imagery and to improve the accuracy of locations recorded from the air. A general sketch of the area and/or photograph will be taken for each use-site. The *Use Site ID* variable connects locations used by each crane group. The *Use Site ID* is a sequential number (beginning with 01) assigned by the project manager and will be recorded on the datasheets with the corresponding *Crane Group ID*, *Location ID* and *Time*.



IV.C. Data Collection from State and Federal Agencies

The report will contain a summary of all whooping crane migrational sightings within Nebraska and specifically the central Platte River corridor as obtained from the FWS, Grand Island Field Office. FWS crane group identification numbers will be recorded in the database to identify individual crane groups that use the study area for multiple days.

IV.D. Additional Metrics

Metrics outlined in Table 1 as well as additional metrics may be included in habitat selection analyses along with the physical characteristics of crane use locations that are recorded by the Consultant. All metrics will be measured remotely via GIS, LiDAR data, aerial imagery, HEC-RAS model, and data collected under other protocols (e.g., Geomorphology and In-channel Vegetation Monitoring Protocol).

Table 1. Additional metrics and measurements that may be included in whooping crane habitat selection analyses that generally will not be recorded or measured by the Consultant.

| Metric | Metric Description | Use Site Data Source | Available Site Data Source |
|---------------------------|--|---------------------------|----------------------------|
| Proportion of Corn | Proportion of landcover surrounding use locations classified as corn | USDA NASS | USDA NASS |
| Proportion of Wet Meadow | Proportion of landcover surrounding use locations classified as wet meadow | PRRIP WM Coverage | PRRIP WM Coverage |
| Proportion of Grassland | Proportion of landcover surrounding use locations classified as grassland | PRRIP WM Coverage | PRRIP WM Coverage |
| Proportion of Forest | Proportion of landcover surrounding use locations classified as forest | PRRIP HA Coverage | PRRIP HA Coverage |
| Unforested Width | Width of river corridor unobstructed by riparian forest | Aerial Photo | Aerial Photo |
| Unvegetated Channel Width | Width of channel clear of dense vegetation | Aerial Photo and/or LiDAR | Aerial Photo and/or LiDAR |
| Total Channel Width | Total width of channel from left bank to right bank (ft) | HEC-RAS model | HEC-RAS model |
| Wetted Width | Top width of wetted channel | HEC-RAS model | HEC-RAS model |
| Proportion Wetted | Proportion of total channel width that is wetted | HEC-RAS model | HEC-RAS model |
| Mean Depth | Mean depth of the wetted portion of the channel (ft) | HEC-RAS model | HEC-RAS model |
| Unit Discharge | Flow (cfs) per linear foot of channel width (Split Flow/Total Channel Width) | HEC-RAS model | HEC-RAS model |
| Width Depth Ratio | Ratio of channel width to depth (Wetted Width / Mean Depth) | HEC-RAS model | HEC-RAS model |

IV.E. Analysis Methods

Information collected through this protocol will be used to evaluate changes in distribution of use and habitat characteristics for whooping crane use-sites in the study area. As such, the Consultant will perform basic data analyses requested by the Program and will report findings that include explanatory as well as graphical representations of their findings on a migrational-season basis and will relate trends in findings such as an increase/decrease in use, increase/decrease in proportion of population stopping in the study area, stop-over duration, etc. to whooping crane population sustainability and growth. This protocol is designed to provide information on crane groups with an estimated probability of inclusion in the sample regardless of location in the study area. Since aerial survey provide this information, but opportunistically



located cranes have an unknown probability of inclusion, separate analyses will be conducted for each of these types of data.

Habitat Use

Since the whooping crane is a rare species and identifying individual cranes is usually not possible, all analyses will need to balance small sample sizes with pseudo-replication. There are two options for analyzing habitat use, one will retain the sample size as the number of individual whooping crane groups, as identified by the FWS, as the first step of the analysis and the second will include every observation of each crane group and will contain multiple observations per group.

There are several analysis methods available for summarizing the habitat characteristics of whooping crane use-sites. The methods range from calculating means and variances, to modeling habitat use, to documenting changes through time, to methods that are not currently developed. With each analysis, the probability-based sample of whooping crane use-sites collected under this protocol will provide data adequate for inferences to all crane stop-over-sights within the central Platte River study area.

Indices of Use

Annual Indices of crane use will be developed using information obtained by this protocol. Indices of use will document the proportion of the population and number of cranes and crane groups observed per survey effort (flight, migration season, etc.) using Program defined and evaluated suitable habitat, Program habitat complexes, and habitat classified as ‘unsuitable’ by existing Program standards. Habitat availability assessments will be conducted annually under a separate Program contract and the results will be available to the Consultant. If the protocol is implemented in a consistent manner, a change in these indices through time will estimate changes in the frequency of use of the study area, Program habitat complexes, as well as Program-defined suitable and unsuitable habitat throughout the first increment.

V. QUALITY ASSURANCE/QUALITY CONTROL (QA/QC)

QA/QC measures will be implemented at all stages of the study, including field data collection, data entry, data analysis, and report preparation. Observers will be trained and tested in the methods used and on their ability to identify whooping cranes. Data forms will be completed on a daily basis. At the end of each survey day, each observer will be responsible for inspecting his or her data forms for completeness, accuracy, and legibility. The study team leader will review data forms to insure completeness and legibility, and correct the forms as needed. Any changes made to the data forms will be initialized by the person making the change.

To help train observers that will be conducting the aerial surveys, each individual will be required to fly practice transects, or a portion of one transect. During this flight there will be whooping crane decoys placed in the river channel to allow observers the opportunity to see a “whooping crane” from the air at the speed and altitude of the surveys.

Data will be entered into the Program’s database by qualified technicians. These files will be compared to the raw data forms and checked for errors. Irregular codes detected or any ambiguous data will be discussed with the observer and study team leader to clarify and document changes.

After data have been keyed and verified, the study team leader or QA/QC technician will check data forms against the final computer file and any problems identified will be traced back to raw data forms, observer(s), and data-entry personnel and corrections will be documented.

VI. DATA COMPILATION AND STORAGE

The Program’s database will be used to store, retrieve, and organize field observations. Microsoft Office InfoPath is required to enter data into the database. The data for each survey will be incorporated within the larger Program database. All field data forms, field notebooks, and electronic data files will be retained for ready reference.



VII. REPORT FORMAT

Data on whooping crane habitat use will be compiled, summarized, and incorporated within the larger Program database following each migration season. Migration-period reports will be submitted to the Program's Executive Director's Office (EDO) for review and distribution following each migration season. Draft and final reports that summarize findings and describes methods, analyses (including descriptive statistics of whooping crane use), results, and any conclusions that can be drawn will have both written and graphical components and will also contain maps and/or aerial photos showing crane use-sites. Reports will be submitted to the Program's EDO for review and distribution to the Technical Advisory and Governance Committees.


VIII. DATA SHEETS – *To be provided prior to survey implementation*

- Aerial Survey
- Aerial Observation



Appendix A. Program Associated Habitat Area.


Cooperative Agreement Platte River Study Area



Maps and Analysis Prepared by Lynx Spatial Technologies Inc.




Appendix B. Riverine and targeted primary and secondary wetland return flight transects that will be flown daily during systematic aerial wetland roost surveys.





Appendix C. Median (dots) and fifth and ninety-fifth percentiles (bars) of initial observation dates of whooping cranes in Nebraska during spring migration. Horizontal solid lines represent the spring monitoring season in the Program's whooping crane monitoring protocol that has been implemented since Spring 2014. Horizontal dashed line represents the start of the spring monitoring season in the Program's whooping crane monitoring protocol that was implemented Spring 2001–Spring 2013. These dates were based on the fifth and ninety-fifth percentiles of initial observation dates of whooping cranes in Nebraska, 1975–1999 (~reproduced in the right plot). Vertical dashed lines separate the 10, 20, and 40 year analyses from analyses of telemetry data (2010–2016) and the 1975–1999 USFWS whooping crane data.





Appendix D. Median (dots) and fifth and ninety-fifth percentiles (bars) of initial observation dates of whooping cranes in Nebraska during fall migration. Horizontal solid lines represent the fall monitoring season in the Program's whooping crane monitoring protocol that has been implemented since Fall 2017. Horizontal dashed line represents the end of the fall monitoring season in the Program's whooping crane monitoring protocol that was implemented Fall 2001–Fall 2016. These dates were based on the fifth and ninety-fifth percentiles of initial observation dates of whooping cranes in Nebraska, 1975–1999 (~reproduced in the right plot). Vertical dashed lines separate the 10, 20, and 40 year analyses from analyses of telemetry data (2010–2016) and the 1975–1999 USFWS whooping crane data.

